

Bell Violation with Discrete-Continuous Entanglement

X.-F. Qian,* C.J. Broadbent, and J.H. Eberly

*Rochester Theory Center and the Department of Physics & Astronomy
University of Rochester, Rochester, New York 14627*

(Dated: February 26, 2013)

We describe a new Bell test for two-party entangled systems, one that engages an unbounded continuous variable. The continuous variable state is allowed to be arbitrary and inaccessible to direct measurements. A systematic method is introduced to perform the required measurements indirectly. Our results provide a new perspective both in the study of local realistic theory for continuous variable systems and in the nonlocal control theory of quantum information.

PACS numbers: 03.65.Ud, 42.50.-p, 42.50.Ex

The issue of incompatibility between local realism and the completeness of quantum mechanics was originally raised for unbounded continuous variables in two-party systems by Einstein, Podolsky, and Rosen (EPR) [1]. Experiments based on inequalities to test local realism proposed by Bell [2] and his followers [3] show, as is well known, that in at least some physical domains quantum theory is fully compatible with their results while classical realism is not. The initial, and most of the subsequent, experimental tests of local realism have been realized without engaging continuous variables but rather using the discrete variables available in photon polarization [4–6], trapped ions [7], solid-state qubits [8], etc.

The major difficulty in extending standard Bell tests to unbounded continuous variable state spaces is the need to identify a finite set of realistic measurement operators that (a) are adequate for systematic examination of a continuous state space, and (b) can be bounded in some fashion that makes possible the derivation of a Bell inequality.

One proposed approach to avoid much of the difficulty has been to discretize the continuous spaces with binning or pseudo-spin methods that reduce the doubly continuous infinity of dimensions to two or a few, in which case one can map onto the familiar analysis of discrete Bell-like inequalities (see, e.g., [9–15]). However, it remains true that even if these methods produce a reduced space that is easy to measure and manipulate, they have treated only a fraction of the continuous variable space and are always open to the question why a particular binning or pseudo-spin scheme was chosen.

A non-dimension-reducing approach by Cavalcanti, Foster, Reid, and Drummond (CFRD) [16] leads to a continuous multipartite inequality that doesn't rely on a discretization of the continuous degrees of freedom. However, to construct the CFRD inequality, operator commutation relations must be ignored, which also eliminates a large category from test (see Sun, et al. [17]). Additionally, violation of these inequalities may not be possible with only two parties [18].

The central issue to be addressed here has two parts: to derive a standard CHSH Bell inequality [3] for an ar-

bitrary bipartite input state in an unbounded continuous variable state space, and to describe a currently accessible experimental method for its test. We provide what we believe is the first step in this direction by treating a two-party discrete-continuum state. One party is described by an arbitrary continuum state and the other party by a discrete two-dimensional state. To provide easy visualization, we address both issues in a specific scenario using the following two-photon down-conversion state:

$$|\psi_{AB}\rangle = \cos\theta |H\rangle_A \otimes \int d\vec{q} h(\vec{q}) |\vec{q}\rangle_B + \sin\theta |V\rangle_A \otimes \int d\vec{q} v(\vec{q}) |\vec{q}\rangle_B, \quad (1)$$

where $|\vec{q}\rangle_B$ is one of a continuum of delta-normalized one-photon transverse momentum states of photon B , and $|H\rangle_A$ and $|V\rangle_A$ denote horizontally and vertically polarized quantum states of photon A . We assume that the transverse momentum state of photon A and the polarization of photon B factor out of the quantum state, and therefore need not be indicated.

The $\sin\theta$ and $\cos\theta$ factors are included in writing $|\psi_{AB}\rangle$ to preserve its unit normalization, as the complex continuum amplitudes $h(\vec{q})$ and $v(\vec{q})$ are assumed to be unit-normalized, i.e., $\int d\vec{q} |h(\vec{q})|^2 = \int d\vec{q} |v(\vec{q})|^2 = 1$. Beyond normalization, nothing else is assumed about $h(\vec{q})$ and $v(\vec{q})$, including the value of their generally non-zero scalar product,

$$\int d\vec{q} h^*(\vec{q}) v(\vec{q}) \equiv z \neq 0. \quad (2)$$

The two-photon state in (1) has an important freedom in the amplitude functions $h(\vec{q})$ and $v(\vec{q})$, which are arbitrary superpositions of the modes in continuous $\vec{q}$ space. In the following we will use the term *bundle* to refer to an arbitrary superposition of $|\vec{q}\rangle$ states. Note that this means that it is impossible to fully determine the state (infinitely many measurements would be required). This point is crucial because it is the stopping point for attempts up to the present time to fully engage a continuous degree of freedom in Bell inequality analysis. We have overcome this roadblock, as we describe below.

It is natural to use the Schmidt analysis in considering two-party pure state entanglement, whether discrete or continuous. The Schmidt decomposition [19] reformulates the state (1) as

$$|\psi_{AB}\rangle = \sum_{n=1}^{\infty} \kappa_n |u_n\rangle_A |f_n\rangle_B, \quad (3)$$

where the sets $\{|u_n\rangle_A\}$ and $\{|f_n\rangle_B\}$ are superpositions of A 's polarization states and B 's momentum states respectively, and are derivable as the eigenvectors of A 's discrete and B 's continuous reduced density matrices. The κ_n^2 are the associated eigenvalues, which are always the same for the two reduced density matrices.

Note that since party A has only two dimensions it has only two eigenvalues, and this forces all but two of B 's infinitely many Schmidt eigenvalues to vanish. Thus the infinite n sum in (3) has only two non-zero terms, which we write:

$$|\psi\rangle = \kappa_1 |u_1\rangle |f_1\rangle + \kappa_2 |u_2\rangle |f_2\rangle, \quad (4)$$

where we have dropped the A and B labels because it will be easy to remember that the discrete states belong to photon A and the continuous states to photon B . Without loss of generality we assume that $\kappa_1 \geq \kappa_2$. Here $|u_1\rangle$ and $|u_2\rangle$ are merely rotations of the original polarization states $|H\rangle$ and $|V\rangle$, and $|f_1\rangle$ and $|f_2\rangle$ are bundles of B 's momentum states $|\vec{q}\rangle$ where the bundles are determined by the Schmidt rearrangement [20], and we write them as $|f_1\rangle = \int d\vec{q} \varphi_1(\vec{q}) |\vec{q}\rangle$ and $|f_2\rangle = \int d\vec{q} \varphi_2(\vec{q}) |\vec{q}\rangle$, with a key orthogonality property:

$$\langle f_1 | f_2 \rangle = \int d\vec{q} \varphi_1^*(\vec{q}) \varphi_2(\vec{q}) = 0, \quad (5)$$

while κ_1 and κ_2 are real positive coefficients analogous to the $\sin \theta$ and $\cos \theta$ in (1), with $\kappa_1^2 + \kappa_2^2 = 1$. We note that because $h(\vec{q})$ and $v(\vec{q})$ are unknown, then $\{\kappa_1, \kappa_2\}$ are also unknown. Lastly, for simplicity in the following derivation, we assume that z is real-valued which ensures that $|u_i\rangle$ is linearly polarized.

The Schmidt theorem provides an ‘‘optimum’’ result in three ways. First, as partners for the rotated polarization states it makes two bundles of momentum states using the (presumed unknown) amplitudes $h(\vec{q})$ and $v(\vec{q})$. Second, it guarantees that those state bundles are orthogonal, and so we have a pair of orthonormality relations $\langle u_i | u_j \rangle = \langle f_i | f_j \rangle = \delta_{ij}$. Third, independent of the makeup of the two bundles, the Schmidt states $|f_1\rangle$ and $|f_2\rangle$ define a plane in the infinite dimensional $|\vec{q}\rangle$ space.

We are now much closer to Bell Inequality territory because rotations in planes in A and B spaces are what the CHSH inequality demands. But the bundles of continuum states making up the two Schmidt states $|f_1\rangle$ and $|f_2\rangle$ are mysterious because the original functions $h(\vec{q})$ and $v(\vec{q})$ were unknown. There are no operators available

in continuum B space to make the rotations required by the CHSH Bell analysis. We will describe below how to make measurements in a rotated basis in the continuum space without rotation operators for the space, but first, let us reproduce the CHSH Bell Inequality analysis, under the assumption that rotations in the $|f_1\rangle$ - $|f_2\rangle$ plane can be controlled.

With ordinary optical components one can always undertake a rotation of the Schmidt basis in photon A 's polarization space, i.e.,

$$|u_1^\alpha\rangle = \cos \alpha |u_1\rangle + \sin \alpha |u_2\rangle, \quad (6)$$

$$|u_2^\alpha\rangle = -\sin \alpha |u_1\rangle + \cos \alpha |u_2\rangle, \quad (7)$$

where α is an arbitrarily adjustable parameter that defines the rotation angle. A rotated basis $|f_1^\beta\rangle, |f_2^\beta\rangle$ of momentum space bundles for photon B can be defined similarly with β as the rotation angle in $\vec{q}$ space, while the practical matter of accomplishing such a rotation remains temporarily an open question.

However, given these rotations, the conventional CHSH analysis of local hidden variable theory [3] can be employed. One considers the Bell operator $\mathcal{B}$ and finds $\mathcal{B} \leq 2$, where $\mathcal{B}$ is defined as

$$\mathcal{B} = C(\alpha, \beta) - C(\alpha, \beta') + C(\alpha', \beta) + C(\alpha', \beta'). \quad (8)$$

Here $C(\alpha, \beta)$ is the CHSH correlation between photons A and B when the measurements are set for the angles α and β , and $P_{ij}(\alpha, \beta)$ are the joint probabilities of finding photon A in state $|u_i^\alpha\rangle$ and photon B in state $|f_j^\beta\rangle$, with $i, j = 1, 2$. That is,

$$C(\alpha, \beta) = P_{11}(\alpha, \beta) - P_{12}(\alpha, \beta) - P_{21}(\alpha, \beta) + P_{22}(\alpha, \beta). \quad (9)$$

According to quantum mechanics, the joint probability is given as $P_{ij}(\alpha, \beta) = \langle \psi_{AB} | u_i^\alpha \rangle \langle f_j^\beta | \langle u_i^\alpha | \psi_{AB} \rangle$, which is a joint projection in the state spaces of both photons and has the potential to violate the CHSH inequality. Then the Bell operator $\mathcal{B}$ can be calculated to be

$$\begin{aligned} \mathcal{B} = & 2\kappa_1 \kappa_2 \left[\sin 2\alpha (\sin 2\beta - \sin 2\beta') \right. \\ & \left. + \sin 2\alpha' (\sin 2\beta + \sin 2\beta') \right] \\ & + \cos 2\alpha (\cos 2\beta - \cos 2\beta') \\ & + \cos 2\alpha' (\cos 2\beta + \cos 2\beta'). \end{aligned} \quad (10)$$

For the choices $\alpha = 0$, $\beta = \pi/8$, $\alpha' = \alpha + \pi/4$ and $\beta' = \beta + \pi/4$, one finds

$$\mathcal{B} = \sqrt{2}(2\kappa_1 \kappa_2 + 1). \quad (11)$$

Then there will be a Bell violation, $\mathcal{B} > 2$, whenever $\kappa_1 \kappa_2 > \sqrt{2} - 1$. Obviously this can be satisfied and for




FIG. 1: Schematic illustration of Bell test for discrete-continuum entangled photon pairs. The red source S_t emits a photon pair $|\psi\rangle_{t\bar{t}}$, where the discrete (polarization) space of the photon propagating towards northwest (NW) in mode t is entangled with the continuous (e.g., momentum) space of the photon heading southeast (SE) in mode $\bar{t}$. The blue source S_a emits identical photon pairs denoted as $|\psi\rangle_{a\bar{a}}$ with the discretely and continuously entangled photons propagating towards southwest (SW) and northeast (NE) respectively. The photon in mode t passes through a polarizer $\hat{u}_t^\alpha$ that passes only the polarization component $|u_1^\alpha\rangle$, and then enters mode T for detection. Similar actions are taken for the photon in mode a with a polarizer $\hat{u}_a^s$ that passes only the polarization component $|u_1^s\rangle$. The photons in modes $\bar{t}$ and $\bar{a}$ are combined by a 50:50 beam splitter (BS) with two outcome modes $\bar{T}$ and $\bar{A}$ being detected.

the state with $\kappa_1 = \kappa_2 = 1/\sqrt{2}$ even the Cirelson bound is attained, i.e., $\mathcal{B}$ reaches the maximum value $2\sqrt{2}$.

As described above, the central hurdle to be overcome is the lack of a method to measure the Schmidt bundles in the continuous $|\vec{q}\rangle$ space of photon B . As we now demonstrate, a specially engineered auxiliary photon is sufficient to accomplish this [21]. The requisite auxiliary photon can be easily created using an auxiliary entangled state which is identical to the original state. Practical techniques for generating pairs of identical entangled biphotons are available, as discussed in the Supplemental Information, so we proceed with the setup sketched in Fig. 1. The two-party state to be Bell-tested is labeled with the subscripts t and $\bar{t}$ for its two photons, and the labels a and $\bar{a}$ are used for the identical auxiliary pair.

The two sources emit pairs of entangled photons propagating along directions t and $\bar{t}$, and a and $\bar{a}$. The Schmidt form of the discrete-continuum state of the entangled photon pair under test is

$$|\psi\rangle_{t\bar{t}} = \kappa_1 |u_1\rangle_t |f_1\rangle_{\bar{t}} + \kappa_2 |u_2\rangle_t |f_2\rangle_{\bar{t}}, \quad (12)$$

and in the figure the discretely (polarization) entangled photon in mode t is heading northwest (NW) and the continuously (momentum) entangled photon in mode $\bar{t}$ is heading southeast (SE), illustrated by the red paths in Fig. 1. The auxiliary photon pair has exactly the same state

$$|\psi\rangle_{a\bar{a}} = \kappa_1 |u_1\rangle_a |f_1\rangle_{\bar{a}} + \kappa_2 |u_2\rangle_a |f_2\rangle_{\bar{a}}, \quad (13)$$

with the discretely entangled photon in mode a heading SW and the continuously entangled photon in mode $\bar{a}$ heading NE, illustrated by the blue paths in Fig. 1.

A polarization projection, e.g., on basis $|u_1^\alpha\rangle$, for the photon in mode t can be realized with a polarizer $\hat{u}_t^\alpha$ that passes the $|u_1^\alpha\rangle$ component into mode T , i.e.,

$$|u_1^\alpha\rangle_t \langle u_1^\alpha| \psi \rangle_{t\bar{t}} = |u_1^\alpha\rangle_T \{ \kappa_1 c_\alpha |f_1\rangle_{\bar{t}} + \kappa_2 s_\alpha |f_2\rangle_{\bar{t}} \} \quad (14)$$

where c_α and s_α stand for $\cos \alpha$ and $\sin \alpha$. The probability of this measurement outcome being realized is given by $P_1(\alpha) = {}_{t\bar{t}}\langle \psi | u_1^\alpha \rangle_t \langle u_1^\alpha | \psi \rangle_{t\bar{t}} = \kappa_1^2 c_\alpha^2 + \kappa_2^2 s_\alpha^2$, and can be determined experimentally by blocking mode $\bar{a}$ and recording the number of coincidences detected during a fixed time window in modes $(T, \bar{T})$ and $(T, \bar{A})$ for angles $\{\alpha, \alpha + \pi/2\}$,

$$P_1(\alpha) = \frac{N_\alpha(T, \bar{T}) + N_\alpha(T, \bar{A})}{\sum_{\delta \in \{\alpha, \alpha + \pi/2\}} (N_\delta(T, \bar{T}) + N_\delta(T, \bar{A}))}, \quad (15)$$

where $N_\delta(T, Y)$ is the number of coincidences in modes T and Y with the polarizer in mode T at angle δ . We note here that the origin of the angle α is not known a priori but can be determined by rotating the polarizer until the output is at a maximum, $P_1(\alpha = 0) = \max_\alpha P_1(\alpha)$. Coincidentally, this also determines κ_1 and κ_2 , since $P_1(0) = \kappa_1^2$ and $\kappa_1^2 + \kappa_2^2 = 1$.

We want to measure the continuum space in a basis rotated by the angle β as well, so we now express the state in the rotated basis, $\{|f_1^\beta\rangle, |f_2^\beta\rangle\}$,

$$|u_1^\alpha\rangle_t \langle u_1^\alpha| \psi \rangle_{t\bar{t}} = |u_1^\alpha\rangle_T \left\{ (\kappa_1 c_\alpha c_\beta + \kappa_2 s_\alpha s_\beta) |f_1^\beta\rangle_{\bar{t}} + (-\kappa_1 c_\alpha s_\beta + \kappa_2 s_\alpha c_\beta) |f_2^\beta\rangle_{\bar{t}} \right\}, \quad (16)$$

which we rewrite again as

$$|u_1^\alpha\rangle_t \langle u_1^\alpha| \psi \rangle_{t\bar{t}} = \sqrt{P_1(\alpha)} |u_1^\alpha\rangle_T \left(c_{11} |f_1^\beta\rangle_{\bar{t}} + c_{12} |f_2^\beta\rangle_{\bar{t}} \right). \quad (17)$$

Here c_{ij} with $i, j = 1, 2$, are normalized amplitude coefficients, and they relate to joint probabilities in an obvious way: $P_{ij}(\alpha, \beta) = |c_{ij}|^2 P_i(\alpha)$.

A projection in the continuous variable space, e.g., on basis $|f_j^\beta\rangle$, for the photon in mode $\bar{t}$ can be realized with the help of the auxiliary photon pair $|\psi\rangle_{a\bar{a}}$. First, the mode a photon of the auxiliary pair is projected (by a polarizer $\hat{u}_a^s$) onto the polarization basis $|u_1^s\rangle$, where angle s is chosen to strip off the $|f_2^\beta\rangle$ component from the photon in mode $\bar{a}$. A glance at (16) shows how a stripping in continuum space by action in polarization space works. In (16), by choosing α such that $\kappa_1 \tan \beta = \kappa_2 \tan \alpha$, the $|f_2\rangle$ component would be eliminated. In the case of auxiliary photon a , we choose s such that $\kappa_1 \tan \beta = \kappa_2 \tan s$ and obtain

$$|u_1^s\rangle_a \langle u_1^s| \psi \rangle_{a\bar{a}} = \sqrt{P_1(s)} |u_1^s\rangle_A |f_1^\beta\rangle_{\bar{a}}, \quad (18)$$

with $P_1(s) = {}_{a\bar{a}}\langle\psi|u_1^s\rangle_a {}_a\langle u_1^s|\psi\rangle_{a\bar{a}}$. $P_j(s)$ is determined experimentally in exactly the same way as $P_i(\alpha)$. The photon enters mode A from mode a after passing the stripping polarizer $\hat{u}_1^s$, as shown in Fig. 1. Then the four-photon state after the two polarization projections in modes t and a is given by

$$|\psi\rangle_{T\bar{t}A\bar{a}} = \sqrt{P_1(\alpha)P_1(s)}|u_1^\alpha\rangle_T|u_1^s\rangle_A \otimes (c_{11}|f_1^\beta\rangle_{\bar{t}} + c_{12}|f_2^\beta\rangle_{\bar{t}}) \otimes |f_1^\beta\rangle_{\bar{a}}. \quad (19)$$

Next, as shown in Fig. 1, the mode $\bar{t}$ photon is combined with the mode $\bar{a}$ photon (which is in the continuous variable state $|f_1^\beta\rangle$) by a 50:50 beam splitter (BS). The outcome modes are denoted as $\bar{T}$ (NE) and $\bar{A}$ (SE). The effect of the BS can be expressed as

$$|f_j^\beta\rangle_{\bar{t}} = (|f_j^\beta\rangle_{\bar{A}} + i|f_j^\beta\rangle_{\bar{T}})/\sqrt{2}, \quad (20)$$

$$|f_j^\beta\rangle_{\bar{a}} = (i|f_j^\beta\rangle_{\bar{A}} + |f_j^\beta\rangle_{\bar{T}})/\sqrt{2}. \quad (21)$$

The component of the four-photon state after the BS which has only a single photon in $\bar{T}$ and $\bar{A}$ is given by

$$|\psi\rangle_{T\bar{T}A\bar{A}} = \sqrt{P_1(\alpha)P_1(s)}|u_1^\alpha\rangle_T|u_1^s\rangle_A c_{12} \otimes (|f_1^\beta\rangle_{\bar{T}}|f_2^\beta\rangle_{\bar{A}} - |f_2^\beta\rangle_{\bar{T}}|f_1^\beta\rangle_{\bar{A}})/2. \quad (22)$$

One notes that due to the Hong-Ou-Mandel (HOM) effect [2] the only contributing component of the mode $\bar{t}$ photon in Eq. (19) to the state after the BS is $|f_2^\beta\rangle_{\bar{t}}$. This amounts to a filtering or projecting operation of the photon in mode $\bar{t}$ into the continuous variable basis $|f_2^\beta\rangle$.

With the above operations, a joint projection is realized for the testing entangled photon pair $|\psi\rangle_{t\bar{t}}$. It is then straightforward to achieve the joint probability $P_{12}(\alpha, \beta)$. The four-photon coincidence probability in modes $T, \bar{T}, A, \bar{A}$ is given as

$$P_{T\bar{T}A\bar{A}}(\alpha, \beta) = \frac{P_1(\alpha)P_1(s)c_{12}^2}{2} = \frac{N_{\alpha\beta}(T, A, \bar{T}, \bar{A})}{N(t, a, \bar{t}, \bar{a})}, \quad (23)$$

where $N_{\alpha\beta}(T, A, \bar{T}, \bar{A})$ and $N(t, a, \bar{t}, \bar{a})$ are four-photon coincidence counts of the corresponding modes. The individual probabilities can be determined using (15). Consequently, the joint probability can be written in terms of measurable quantities,

$$P_{12}(\alpha, \beta) = \frac{2}{P_1(s)} \frac{N_{\alpha\beta}(T, A, \bar{T}, \bar{A})}{N(t, a, \bar{t}, \bar{a})}. \quad (24)$$

Measurement of the other joint probabilities $P_{11}(\alpha, \beta)$ and $P_{2j}(\alpha, \beta)$ are accomplished by appropriately rotating the angles α and β by $\pi/2$. In this way the correlation function $C(\alpha, \beta)$ can be achieved straightforwardly. Other correlations can be obtained similarly with other choices of angles α and β . To achieve the Bell violation given in (11) the orientation of the stripping polarizer is determined as $\tan s = (\kappa_1/\kappa_2)(\sqrt{2} - 1)$ and $\tan s' = (\kappa_1/\kappa_2)(\sqrt{2} + 1)$.

In summary, we have subjected the CHSH Bell inequality to a new test. We have demonstrated that violations are predicted even when one party is defined by an unbounded continuous variable in a state of unknown complexity. This means that we believe we have treated for the first time the generic case when the continuous state to be tested is not subjected to any binned outcome of detection and is not known to be an eigenstate of any operator or finite set of operators. Thus no operators on the continuous variable are available to generate the continuum of rotations embodied in the CHSH inequality. In addition, no measurement-oriented dimension reduction, which may reduce the original entanglement, is required in our method. This opens a new prospect for studying local realism tests for continuous variable systems.

Our method of measuring the continuous variable space is an example of non-local control (see [23]). It provides a new perspective on indirect measurement of a system state which is not easily accessible experimentally. We have shown explicitly how manipulating its discrete and controllable entangled partner, arbitrary measurements of a system may be made. Apart from increased measurement capabilities, this type of indirect measurement may be useful for transferring or encoding information into continuous variable spaces which are difficult to detect or probe directly. Therefore, with proper design, it may be possible to construct communication protocols which impede potential eavesdroppers from obtaining the encoded information. This will be addressed in a future work.

We acknowledge helpful discussions with W.P. Grice, and acknowledge partial financial support from the following agencies: DARPA HR0011-09-1-0008, ARO W911NF-09-1-0385, NSF PHY-0855701.

* Electronic address: xfqian@pas.rochester.edu

- [1] A. Einstein, B. Podolsky, and N. Rosen, Phys. Rev. Lett. **47**, 777 (1935).
- [2] J.S. Bell, Physics **1**, 195-200 (1964).
- [3] J.F. Clauser, M.A. Horne, A. Shimony and R.A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
- [4] S.J. Freedman and J.F. Clauser, Phys. Rev. Lett. **28**, 938 (1972).
- [5] E.S. Fry and R.C. Thompson, Phys. Rev. Lett. **37**, 465 (1976).
- [6] A. Aspect, P. Grangier and G. Roger, Phys. Rev. Lett. **49**, 91 (1982), and A. Aspect, J. Dalibard and G. Roger, Phys. Rev. Lett. **49**, 1804 (1982).
- [7] M.A. Rowe, D. Kielpinski, V. Meyer, C.A. Sackett, W.M. Itano, C. Monroe, and D.J. Wineland, Nature **409**, 791 (2001).
- [8] M. Ansmann, H. Wang, R.C. Bialczak, M. Hofheinz, E. Lucero, M. Neeley, A.D. O'Connell, D. Sank, M. Weides, J. Wenner, A.N. Cleland, and J.M. Martinis, Nature **461**, 504 (2009).
- [9] A. Gilchrist, P. Deuar, and M.D. Reid, Phys. Rev. Lett.

- 80, 3169 (1998).
- [10] W.J. Munro and G.J. Milburn, Phys. Rev. Lett. **81**, 4285 (1998); W.J. Munro, Phys. Rev. A **59**, 4197 (1999).
 - [11] Z.B. Chen, J.W. Pan, G. Hou, and Y.D. Zhang, Phys. Rev. Lett. **88**, 040406 (2002).
 - [12] J. Wenger, M. Hafezi, F. Grosshans, R. Tualle-Broui, and P. Grangier, Phys. Rev. A **67**, 012105 (2003).
 - [13] C. Brukner, M.S. Kim, J.-W. Pan, and A. Zeilinger, Phys. Rev. A **68**, 062105 (2003).
 - [14] T. Yarnall, A.F. Abouraddy, B.E.A. Saleh, and M.C. Teich, Phys. Rev. Lett. **99**, 170408 (2007).
 - [15] K.H. Kagalwala, G. DiGiuseppe, A.F. Abouraddy and B.E. Saleh, Nature Phot. **7**, 72-78 (2012).
 - [16] E.G. Cavalcanti, C.J. Foster, M.D. Reid, and P.D. Drummond, Phys. Rev. Lett. **99**, 210405 (2007).
 - [17] Q. Sun, H. Nha, and M.S. Zubairy, PRA **80**, 020101(R) (2009).
 - [18] A. Salles and D. Cavalcanti and A. Acín, Phys. Rev. Lett. **101**, 040404 (2008). Their proof can be extended to arbitrary operators having a scalar commutator by using the Heisenberg uncertainty principle in sum form, in the spirit of Ref. [16].
 - [19] E. Schmidt, Math. Ann. **63**, 433 (1907). See also A. Ekert and P.L. Knight, Am. J. Phys. **63**, 415 (1995) and J.H. Eberly, Laser Phys. **16**, 921 (2006).
 - [20] A similar but fully classical situation has been examined in X.-F. Qian and J.H. Eberly, Opt. Lett. **36**, 4110 (2011).
 - [21] A classical entanglement analog has been worked out by X.-F. Qian and J.H. Eberly (in preparation).
 - [22] C.K. Hong, Z.Y. Ou, and L. Mandel, Phys. Rev. Lett. **59**, 2044 (1987).
 - [23] See, for example, J. Eisert, K. Jacobs, P. Papadopoulos, and M.B. Plenio, Phys. Rev. A **62**, 052317 (2000); D. Collins, N. Linden, and S. Popescu, Phys. Rev. A **64**, 032302 (2001). In fact, the setup in Fig. 1 is essentially the same as that used by M. Pavićić and J. Summhammer, Phys. Rev. Lett. **73**, 3191 (1994), in an early entanglement swapping experiment.

Appendix

At least two approaches are open for generating a discrete-continuum (e.g., polarization-spectrum) entangled state to perform the Bell test proposed in the text. One setup is illustrated in Fig. 2, where the entangled photon pairs are produced in a pair of spontaneous parametric down-conversion (SPDC) crystals, a combination of type I and type II, pumped with an ultra-short UV laser pulse in a double pass configuration [1]. The photon pair produced by the first passage can be written in general as

$$|\psi'\rangle_{t\bar{t}} = \int \int d\omega_1 d\omega_2 \Phi(\omega_1, \omega_2) |\omega_1, H\rangle_t |\omega_2, H\rangle_{\bar{t}} + \int \int d\omega_1 d\omega_2 \Psi(\omega_1, \omega_2) |\omega_1, V\rangle_t |\omega_2, H\rangle_{\bar{t}} \quad (25)$$

where $\Phi(\omega_1, \omega_2)$, $\Psi(\omega_1, \omega_2)$ are two different amplitude functions relating to the field-crystal interaction parameters, and ω_1, ω_2 represent the frequency of the signal

(mode t) and idler (mode $\bar{t}$) photons respectively. Here $|\omega_1, H\rangle$ represents a single photon state with frequency ω_1 and polarization H . The first and second terms in Eq. (25) are generated by the type I and type II crystals, respectively. The propagation directions of the two down-converted photons are determined by the phase-matching conditions of the SPDC crystals.




FIG. 2: Schematic illustration of producing two identical discrete-continuum entangled photon pairs with two-pass spontaneous parametric down conversion. The ultra-short UV laser pulse passes through the combination of type I and type II crystals and creates the first photon pair $|\psi'\rangle_{t\bar{t}}$ (in red paths) with the signal and idler photons propagating in modes t and $\bar{t}$ respectively. The UV pulse is then reflected back by the mirror M and passes through the two-crystal structure again to create the second entangled photon pair $|\psi'\rangle_{a\bar{a}}$ (in blue paths) with the signal and idler photons propagating in modes a and $\bar{a}$ respectively. The spectrum of the signal photons in modes t and a are filtered by the interference filters (IF) so that the desired discrete-continuum (polarization-spectrum) entangled states $|\psi\rangle_{t\bar{t}}$ and $|\psi\rangle_{a\bar{a}}$ are achieved respectively.

Then one can insert an interference filter (IF) centered at ω_0 in front of the signal photon (as shown in Fig. 2). After the filter the two-photon state is left in a desired discrete-continuum (polarization-spectrum) entangled state, i.e.,

$$|\psi\rangle_{t\bar{t}} = \int d\omega_2 [\Phi_{\omega_0}(\omega_2) |H\rangle_t |\omega_2\rangle_{\bar{t}} + \Psi_{\omega_0}(\omega_2) |V\rangle_t |\omega_2\rangle_{\bar{t}}], \quad (26)$$

where $\Phi_{\omega_0}(\omega_2) = \int d\omega_1 f^*(\omega_1 - \omega_0) \Phi(\omega_1, \omega_2)$, $\Psi_{\omega_0}(\omega_2) = \int d\omega_1 f^*(\omega_1 - \omega_0) \Psi(\omega_1, \omega_2)$, and $f(\omega_1 - \omega_0)$ is the spectral response function of the filter. Here we have omitted the factorable components, i.e., the spectral state of the photon in mode t and the polarization state of photon in mode $\bar{t}$.

After the first passage, the UV laser pulse is reflected back by a mirror (M) and then passes through the two-crystal structure again to create the second desired discrete-continuum entangled photon pair $|\psi\rangle_{a\bar{a}}$. As shown in Fig. 2, the two down-converted photons propagate in blue paths with the signal photon in mode a and the idler photon in mode $\bar{a}$. Again the spectrum of the signal photon is filtered by an identical IF centered at ω_0 .

Then the two photons in mode $\bar{t}$ and $\bar{a}$ can be combined by a 50:50 beam splitter as proposed in the text to perform the Bell test measurement. To ensure the temporal indistinguishability of the two photons arriving at the beam splitter, one needs to make sure that the laser pulse length is much shorter than the coherence time of the down-converted photon [1]. By adjusting the distance between the two-crystal structure and the mirror one can achieve the Hong-Ou-Mandel effect [2], and thus realize the necessary temporal indistinguishability.

Another approach for realizing the necessary temporal indistinguishability is to produce SPDC photon pairs with very long coherence times by using a very narrow-band filter as demonstrated in Ref. [3]. The temporal indistinguishability is then provided by appropriate post-

selection of coincident detection events in fast single-photon detectors. In this case the second (auxiliary) discrete-continuum entangled pair can be generated from an identical yet independent two-crystal structure.

* Electronic address: xfqian@pas.rochester.edu

- [1] D. Bouwmeester, J.-W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, *Nature* **390**, 575 (1997).
- [2] C. K. Hong, Z. Y. Ou, and L. Mandel, *Phys. Rev. Lett.* **59**, 2044 (1987).
- [3] M. Halder, A. Beveratos, N. Gisin, V. Scarani, C. Simon, and H. Zbinden, *Nature Phys.* **3**, 692 (2007).